# **EXHIBIT 44**

3666

CHPMICAL SYNTHESIS OF MIDIFIED OLICONUCLECTIDES AND THEIR UTILITY AS NON-RADIOACTIVE HYBRIDIZATION PROBES. Jerry L. Ruth® and Robert N. Bryan®, (SFON:N. KAPLAN), Molecular Blosystems, Inc., San Diego, CA 92121

Several decognitions analogs with a primary amine "linker arm" of eight or twelve atoms in length attached at C-5 (1) have been efficiently prepared. These linker arm nucleosides ("LANS") are appropriately blocked and chemically incorporated into oligonucleotides in good yields. This method allows total control over extent and site of modification in the oligomer.

1:R<sub>1</sub> = H R<sub>2</sub> = decayribose 2:R<sub>2</sub> = Oligorucleotide alR<sub>1</sub> = H blR<sub>1</sub> = fluorescein clR<sub>1</sub> = caproylbiotin

Using a modified phosphochloridite method, three Ziners complementary to known sequences of herpes simplex virus (HSV) were constructed to be specific for either HSV-1, HSV-2, or both (generic). The purified oligomers were kinased and hybridized to HSV target DNA. Targets included an HSV-containing plasmid, lab strains of HSV, and clinical samples of both types 1 and 2. Results clearly demonstrated the desired selectivity. Oligonucleotides containing one or more LANs were selectively modified by attachment of fluorescein (2b) or biotin (2c). Biotinylated probes (2c) kinased and hybridized normally. Non-radioactive detection of 2c was accomplished by dye formation using an avidin-dehydrogenase complex. Results indicate such modified oligonucleotides can: 1) be made very efficiently by chemical methods in large amounts; 2) detect target DNA cleanly and specifically; and 3) be detected by non-radioactive methods.

300

SYNTHESIS, THERMODYNAMICS AND DRUG INTERACTIONS OF MODEL OCTANUCLEOTIDES. I. Lassalle\*, H.L. Weith and S.R. Byrn\*. Purgue University, West Lafayette, Indiana 47907.

Two pairs of complementary octanucleotides were chemically synthesized using the modified phosphotriester method. Their sequences are: TGACGTGA - TCACGTCA and TGAGGTGA - TCACCTCA. Optical studies were performed on the single strands in order to accurately determine their molar absorptivity coefficient (c) and their % hyperchromicity. The thermodynamics of helix formation were determined from melting curve data. In addition, the binding of the acridine AMSA to both the single strands as well as the duplexes was investigated. This data provides insight into the parameters controlling helix formation and drug binding.

Sponsored in part by grants CA30234 and GM29175.

## FILE COPY

3668

LAMBDA PHAGE Cro REPRESSOR-DNA INTERACTIONS: PLUOROURACIL ANALOGUES OF OR . William T. Metzler\*, Elizabeth Tecza\*, Kim T. Arndt\* and Ponzy Lu. Department of Chemistry, Univ. of Pennsylvania, Philadelphia, PA 19104

We have synthesized 15 base pair analogues of the center of the lambda  $0_{\rm R}3$  DNA sequence which contain deoxyfluorouracil replacements at specific thymine locations, 5 and 10, in the sequence below [indicated by \* using the numbering system of Kawashima et al. (1977), Biochemistry 16, 4209-4216]:

2 5 10 15 ATCACCGCAAGGGAT TAGTGGCGTTCCCTA

Using quenching of the fluoresence at the tyrosines of the cro repressor molecule, we were able to demonstrate that the fluoroanalogue of Og3 binds to the protein as well as a 15 base pair non fluorine containing Og3 sequence.

pair non fluorine containing  $O_R3$  sequence. By observing the Fluorine-19 NMR signals in the presence and absence of the  $\underline{cro}$  repressor we are able to monitor the formation of the specific  $\underline{cro}$ - $O_R3$  complex. (Supported by grants from the NIH.)

3670

NEW METHOD FOR SEQUENCE ANALYSIS OF OLIGODEOXYRIBONUCLEOTIDES.

D. M. Black\* and P. T. Gilham. Purdue Univ., West Lafayette,
TN 47907

A simple acquencing procedure employs a two-dimensional separation on a single thin layer chromatographic sheet and consists of, in the first dimension, the fractionation by chain length of a nested set of fragments derived from the oligodeoxyribonuclectide by partial exonuclease action and labelled at their non-common ends, followed by in situ enzymatic degradation of the members of the set and separation of the resulting labelled mononuclectides in the second dimension. Spleen phosphodiesterase and alkaline phosphatase are used to produce the nested set whose members have a common 3' terminus, and these are radioactively labelled at their non-common ends with polynuclectide kinase. The fragments are separated on a polyethyleneimine-cellulose thin layer sheet, using a modification of the solvent system developed by Randerath et al. [Nucleic Acids Res. 1, 1121 (1974)]. The oligomers are them digested in situ to 5' mononuclectides using nuclease Bal 31, nuclease 51, or snake venom phosphodiesterase, and the labelled monomer formed at each position is identified by chromatography in the second dimension with 0.2 M NaOAc adjusted to pH 4.2 with AcOH. This approach to sequence analysis has an added advantage in permitting assignment of the identity and location of any modified or unusual base within the oligonuclectide. (Supported by CA 30234)

3669

A NOVEL UNIVERSAL SUPPORT FOR DNA & RNA SINTHESIS. L. J. Arnold, Jr.\*, R. Lohrmann\*, and J. L. Ruth\*(SPON: N. O. Kaplan), Molecular Biosystems, Inc., San Diego, CA 92121

Current methods of DNA and RNA synthesis have been greatly facilitated by the use of solid supports. Typically such supports have either a silica or organic matrix to which is attached the first nucleoside through a 3' ester-linkage. Since the nucleoside can be any of four ribo- or decoxynthonucleosides, eight such supports are necessary to carry out RNA and DNA synthesis. We have designed a universal support which uses a ribose as a primer with

NUC-0-P-0 H H H B:# the 2' and 3' OH protected by substituted acetyl groups. The support matrix is a organic graft which is non-swellable, is stable to acid and base, and has superior flow properties. DNA or RNA synthesis is carried out in either the 3' or 5'

direction. Once synthesis is complete, the ribose acetyl groups are removed with mild base followed by the selective oxidation of the cis diol by periodate and a \$\textit{\textit{e}}\$-elimination of the formed oligomer. This support has many advantages which include: 1) It is equally useful for RNA or DNA synthesis; 2) Chain elongation may go in either the 3' or 5' direction; 3) The synthesized oligomer may be base and phosphate deblocked before or after removal of the oligomer from the support; 4) The product oligomers are of higher purity since blooding groups can be removed before removal of the oligomer; and 5) After removal of protecting groups the oligomer may be left attached which permits its use as an affinity hybridization support. The functional properties of this support have been confirmed using "t-thymidine labels. Under deblooding conditions, less than 6% of the oligomer is removed. During selective cleavage greater than 80% of the oligomer is eliminated. Using this support, we have synthesized a range of 14-20 mers.

3671

NON RADIOACTIVE BIOTIN - DEPENDENT HYBRIDIZATION/DETECTION USING UNLABELED PROBE DNA. BY: CHRISTINE L. BRAKEL\*, KATHERINE MARKARIAM, AND DEAN L. ENGLEHARDT. ENZÖ BIOCHEM, INC. NEW YORK, NEW YORK 10013.

We have developed a method for the non-radioactive detection of DNA probes that do not contain modified nucleotides. In this system, the probe DNA is 3' terminally labeled with TTP or dATP in the presence of terminal transferase to result in a probe which contains single stranded homopolymeric terminii. After this probe has been hybridized to target DNA, the hybridized molecules are complexed with a complementary homopolymer containing biotinylated nucleotides. Hybridization is visualized using a colormetric detection system for biotin composed of streptavidin (a biotin-binding protein) and a biotinylated enzyme (either acid phosphatase or horseradish peroxidase). Addition of the appropriate substrate yields a colored precipitate. The detection sensitivity of this system was found to be greater than the sensitivity obtained using directly biotinylated probes, prepared either by nick translation or terminal labeling. This new method can be used for the detection of specific sequences on dot blots and Southern transfers.

American Society of Biological Chemists
5th Annual Meeting

The American Association of Immunologists 8th Annual Meeting

St. Louis, Missouri une 3-7, 1984

> ABSTRACTS OF PAPERS 2477-3780 INDEXES

### **Publications Committee**

F. STEPHEN VOGEL, Chairman
JOHN S. COOK
RICHARD M. FORBES
FRANKLIN P. INMAN
HENRY KAMIN
RICHARD McINTOSH
JOSEPH J. McPHILLIPS
ROBERT W. KRAUSS, ex officio
HAROLD F. HARDMAN, ex officio

### **Editorial Board**

L. ROSS HACKLER, Chairman MARK L. ENTMAN DONALD M. MARCUS EMANUEL RUBIN ALAN N. SCHECHTER ROBERT E. STITZEL

### **Executive Editor**

KARL F. HEUMANN 301-530-7100

### **Assistant Executive Editor**

JUDITH B. GANDY 301-530-7119

### Copy Editor

KENDALL SITES 301-530-7108

# Federation Proceedings

Official Publication of the FEDERATION OF AMERICAN SOCIETIE FOR EXPERIMENTAL BIOLOGY

### FUTURE FASEB AND SOCIETY MEETINGS

ASBC/AAI—JUNE 3-7, 1984, St. Louis, MCAPS—JULY 29-AUGUST 3, 1984, Lexington, KASPET—AUGUST 19-23, 1984, Indianapolis, INASCB—NOVEMBER 13-17, 1984, Kansas City, MOFASEB—APRIL 21-26, 1985, Anaheim, CAPASEB—APRIL 21-26, 1985, Anaheim, APRIL 21-26, APRIL

FEDERATION PROCEEDINGS is published 15 times a year, monthly except 2 issues in March and 3 in May, by FASEB. All issues except March (nos. 3 and 4) and May 1 and 4 (nos. 6 and 7) contain papers presented at symposiums and articles especially written for FP. The March issues contain the abstracts of papers submitted for presentation at scientific sessions at the Annual Meeting of the Federation. The May 1 and 4 issues (nos. 6 and 7) contain the abstracts of papers submitted for presentation at scientific sessions of the Annual Meeting of the American Society of Biological Chemists and The American Association of Immunologists. The views expressed in articles and in meeting abstracts are those of the authors and not necessarily those of the Federation. Manuscripts that are considered to be of interest to readers of FP are welcome. All manuscripts submitted (including FASEB or constituent Society symposiums as well as any others submitted) are subject to review and approval by the Editors before publication in FP. Rejected manuscripts may be reconsidered if appealed through the Editorial Board. Second-class postage paid at Bethesda, Maryland, and at additional mailing offices. Copyright © 1984 by the Federation of American Societies for Experimental Biology, 9650 Rockville Pike, Bethesda, Maryland 20814. ISSN 0014-9446. Printed at Lancaster Press, Lancaster, Pennsylvania. All rights reserved. Requests for any reproduction of copyrighted material except first page of a regular article should be made in writing to the Executive Editor at 9650 Rockville Pike, Bethesda, Maryland 20814, and should include an explicit statement of intended use and detailed specification of the material to be reproduced. Telephone 301/530-7119. COPYRIGHT: The code at the bottom of the first page of an article indicates the copyright owner's consent that copies of the article may be made provided that the stated fee is paid through the Copyright Clearance Center, Inc., 21 Congress Street, Salem, Massachusetts 01970.

Business Manager: All business matters should be directed to Mr. Austin H. Henry, Business Manager, FASEB, 9650 Rockville Pike, Bethesda, MD 20814 (301/530-7066).

Advertising Representative: From Steven K. Herlitz, Inc., Steven K. Herlitz, Inc., 404 Park Avenue South, 9th Floor, New York, NY 10016 (212/532-9400).

Advertising Production Coordinator: Stella Koerber, in care of Steven K. Herlitz, Inc., 404 Park Avenue South, 9th Floor, New York, NY 10016 (212/532-9400).

Subscription Price: Members of Constituent Societies \$12.00 per year; Nonmembers (Personal) \$65.00 per year; Institutional \$85.00 per year; Students \$35.00 per year with certification. Postage Differentials: Other Countries + \$20.00. All subscriptions entered on a calendar year basis only and payable in advance. Single Issues: \$12.00, except March 1 and 5, Nos. 3 and 4 (Abstracts), \$35.00 and May 1 and 4, Nos. 6 and 7 (Abstracts), \$35.00. Subscriptions and orders should be sent to: FEDERATION PROCEEDINGS, Subscription Department, 9650 Rockville Pike, Bethesda, Maryland 20814 (301/530-7027).

# -Contents

### BSTRACTS

		•		
1ay 1,	Natural killer cells (nontumor	1000	Basophils, mast cells and IgE	1972
	•	1906	abnormalities	1974
<b>1</b>	effector function	1908	Membrane-bound enzymes in	
	and isotype expression	1909		1975
	Immunoglobulin receptors,		Translational control in	
		1914	eukaryotes	1976
			The secretory pathway	1978
1847		1918	Cell surface recognition	1978
			Gene structure and expression	
1849	mediators of inflammation	1924	in plants	1979
	Effect of lymphokines,		•	
	antibodies and			
1851	immunotherapeutic agents		WEDNESDAY AFTERNOON	i
1853	on tumors	1928	June 6, 1984	•
	Basophils, mast cells and IgE	1933	,	
1856			Peptide growth factors and	
1859		1935		1979
1860				10.0
1863		1937	,	
1867			THURSDAY AFTERNOON	
			June 7, 1984	
	Protein kinase III		3	
1869			Transplantation immunity and	
				1980
				1900
			•	1986
		1330		1991
		1061		1991
		1901		1005
		1064		1995
	grycopi otems	1904		1998
1000				1999
1890	THURSDAY AFTERNOON			2004
				000=
	June 7, 1904			2007
	Calladan in a said	1005		
1300	Cellular receptors for	1967	Nonmuscle contractile and	2012
	immunoglobulin	1969	cytoskeletal systems II	2015
	Effect of lymphokines,			2017
	antibodies and		Protein processing and	
	immunotherapeutic agents			2021
1904		1970		2025
	1841 1843 1847 1849 1851 1853 1856 1859 1860 1863 1867 1879 1874 1879 1881 1884 1886 1887 1888 1890 1892 1896 1900	targets)  Cloned lymphoid cells with effector function Regulation of B cell function and isotype expression Immunoglobulin receptors, lymphocyte markers 1843 Disorders of immune regulation in humans Cellular and humoral mediators of inflammation Effect of lymphokines, antibodies and immunotherapeutic agents on tumors Basophils, mast cells and IgE Clinical disorders of immediate hypersensitivity 1850 Viruses and immunological abnormalities 1861 Biochemical pharmacology II Blood coagulation II Protein kinase III Protein kinase III 1872 Action of toxins 1874 Fibrinolysis	targets)	targets) 1906 Cloned lymphoid cells with effector function 1908 Regulation of B cell function and isotype expression 1909 Immunoglobulin receptors, 1841 1843 Disorders of immune regulation in humans 1918 Cellular and humoral mediators of inflammation 2 Effect of lymphokines, antibodies and immunotherapeutic agents 1855 On tumors 1928 Basophils, mast cells and IgE 1835 Clinical disorders of immediate hypersensitivity 1860 Viruses and immunological abnormalities 1935 Clinical disorders of immediate hypersensitivity 1940 Protein kinase III 1947 Right Proteolytic enzymes and inhibitors 1957 Right Proteolytic enzymes and 1886 Cell surface recognition Gene structure and expression in plants WEDNESDAY AFTERNOON June 7, 1984 THURSDAY AFTERNOON 1944 Protein kinase III 1950 Right Proteolytic enzymes and inhibitors 1951 Right Proteolytic enzymes and 1886 Cell surface proteins and 1967 Cellular interactions 1967 Cellular receptors for immunoglobulin 1969 Effect of lymphokines, antibodies and immunotherapeutic agents 1960 Effect of lymphokines, antibodies and immunotherapeutic agents 1960  Cellular receptors for immunoglobulin 1960 Effect of lymphokines, antibodies and immunotherapeutic agents 1960  Conzymes mechanism of action III Enzyme-multienzyme complexes II Protein synthesis: mechanism Protein processing and turnover II 1970  Protein processing and turnover II 1960 Protein processing and tu

Chromatin and nuclear		Endocytosis and receptor-		Flavoproteins	20
proteins	2029	mediated uptake	2043	Histones, chr matin-associated	1
Cytochrom P-450 III:		DNA and RNA: chemistry	2047	proteins and poly(ADP-	1
structure and function	2032	Eukaryotic gene expression III	2050	ribose)	20
Glycoproteins III	2036	Transport III	2053	Proteolytic enzymes II	20
Fatty acid metabolism II	2040	Metalloproteins	2057	Additional abstract	20

### INDEXES

Author index												2068
Permuted title	index											2088

京都というないは、大学のは、大学の一般の一般である。

The abstracts on pages 1841-2067 were prepared by the authors and then printed by photo-offset without change. Abstracts are not subject to scientific review; therefore, the scientific validity of the results reported is the responsibility of the authors and sponsors. Accuracy, form of citation, designation of materials, acknowledgment of all co-authors and of grant support, terminology, nomenclature, and the like, all remain again the responsibility of the authors and sponsors. Readers should note that the appearance of an abstract in this issue does not necessarily imply future publication of a regular scientific paper.

Asterisk (\*) indicates nonmember of Society to which an abstract was submitted.

It is our pleasure to record the help of the staff of BIOSIS in preparing the indexes, particularly from Ms. Carolyn Marconi and Mr. John R. Thomas.